

**Table 1. 2015 foliar fungicide efficacy trial on Jagalene wheat in College Station, TX.**

Treatment	Rate	Application Timing‡	Company	Yield (bu/a)	Test Wt (lb/bu)	Leaf Rust (Flag Leaf)†
Trivapro	4 oz/a					
Quilt Xcel	10.5 oz/a	Flag Leaf	Syngenta	13.8*	55.1	2.3
Induce	0.125 % v/v					
Approach + Approach Prima	3 oz/a + 6.8 oz/a	Jointing + Flag Leaf	Dupont	13.6	55.6	2.3
Induce	0.25% v/v					
Trivapro	4 oz/a					
Quilt Xcel	10.5 oz/a	Flag Leaf	Syngenta	13.5	55.6	2.3
Agri-Dex (COC)	1 % v/v					
Prosaro 421	5 oz/a	Flag Leaf	Bayer	13.4	55.5	3.0
Induce	.25% v/v					
Custodia	8.6 oz/a	Flag Leaf	MANA	12.4	56.0	3.0
Induce	0.125 % v/v					
Quilt Xcel	10.5 oz/a	Flag Leaf	Syngenta	12.1	55.8	3.0
Induce	0.125 % v/v					
Twinline	9 oz/a	Flag Leaf	BASF	12.1	55.4	1.7
Induce	0.125 % v/v					
Quadris	4 oz/a	Flag Leaf	Syngenta	12.0	55.7	4.3
Induce	0.125 % v/v					
Priaxor	4 oz/a	Flag Leaf	BASF	11.8	55.1	5.0
Induce	0.125 % v/v					
Tebustar 3.6L + Caramba	4 oz/a + 17 oz/a	Jointing + Flag Leaf	Agri Star + BASF	11.6	54.9	3.7
Induce	0.1 % v/v					
Approach Prima	6.8 oz/a	Flag Leaf	Dupont	11.4	55.7	2.3
Induce	0.25% v/v					
Fortix	5 oz/a	Flag Leaf	Cheminova	11.1	55.4	3.7
Induce	0.125 % v/v					
Absolute 500	4 oz/a	Flag Leaf	Bayer	10.6	56.1	4.3
Induce	0.25% v/v					
Tebustar 3.6L (split applic.)	2 oz/a + 2 oz/a	Jointing + Flag Leaf	Agri Star	9.4	53.8	4.3
Induce	0.1 % v/v					
Stratego YLD	4 oz/a	Flag Leaf	Bayer	9.1	54.8	7.0
Induce	0.25% v/v					
Tebustar 3.6L	4 oz/a	Flag Leaf	Agri Star	8.1	55.9	3.7
Induce	0.1 % v/v					
Untreated Control				5.4	51.8	7.0
<b>LSD (P=0.05)</b>				<b>3.3</b>	<b>1.1</b>	<b>2.6</b>
<b>Grand Mean</b>				<b>11.3</b>	<b>55.2</b>	<b>3.7</b>
<b>CV</b>				<b>17.5</b>	<b>1.2</b>	<b>42.6</b>

\*Values in red indicate products with highest statistical ( $P < 0.05$ ) yield, test weight, or disease suppression.

†Leaf rust rating taken on a 0-9 scale (9=100% infection) on 4/21/2015.

‡Foliar fungicide applications were made on 3/6/15 and 4/8/15 for jointing and flag leaf applications, respectively.